

# **1/12° Pacific HYCOM Results**

**E. Joseph Metzger<sup>†</sup>, Harley E. Hurlburt<sup>†</sup>,  
Alan J. Wallcraft<sup>†</sup> and Luis Zamudio<sup>‡</sup>**

**<sup>†</sup>Naval Research Laboratory, Stennis Space Center, MS**

**<sup>‡</sup>Center for Ocean-Atmospheric Prediction Studies, Florida State University**


# Pacific HYCOM Model Configuration

- **Horizontal grid: 1/12° equatorial resolution (2294 x 1362 grid points, 6.5 km spacing on average)**
- **20°S to 65.8°N**
- **20 vertical coordinates**
- **Bathymetry: Quality controlled ETOP05**
- **Surface forcing: (wind stress, wind speed, heat flux [using bulk formula], E-P + relaxation to climatological SSS)**
- **River runoff**
- **Buffer zone: ~3° band along southern and eastern boundary with relaxation to monthly climatological T and S**
- **Closed boundaries along 20°S, in the Indonesian throughflow region and in the Bering Strait**

# **1/12° Pacific HYCOM Modeling Progress**

- **Four 1/12° simulations**
  - **high frequency Hellerman and Rosenstein (1983, JPO) (HR) climatological forced simulation (9.5 years)**
  - **high frequency European Centre for Medium-range Weather Forecasts (ECMWF) climatological forced simulation (8.5 years)**
  - **high frequency ECMWF climatological forced simulation with modification to winds over Hawaii (4 years)**
  - **FNMOC NOGAPS/HR interannual simulation January 2001 – May 2002, a period that spanned the life cycle of Hurricane Juliette**

# 1/12° Pacific HYCOM Basin-scale Circulation SSH Snapshot – 1 January




Forced with high frequency climatological **ECMWF** winds and thermal forcing

# 1/12° Pacific HYCOM

## Zoom on the Kuroshio


### SSH and SST Snapshot – 1 January




Forced with high frequency climatological **ECMWF** winds and thermal forcing

# Comparison of the Basin-scale Circulation MODAS climatology vs. 1/12° Pacific HYCOM

Mean dynamic  
height (dyn cm)  
wrt 1000 db




6-yr mean  
SSH (cm)




Forced with high frequency climatological HR winds and ECMWF thermal forcing

# Comparison of the Basin-scale Circulation MODAS climatology vs. 1/12° Pacific HYCOM

Mean dynamic  
height (dyn cm)  
wrt 1000 db




6-yr mean  
SSH (cm)




Forced with high frequency climatological ECMWF winds and thermal forcing

# 1/12° Pacific HYCOM 6 Year Mean SSH – Kuroshio sub region




**ECMWF forcing**





**HR forcing**

# **1/12° Pacific HYCOM Basin-scale SST 6 year mean**




**Forced with high frequency climatological HR winds and ECMWF thermal forcing**

# Comparison of the Basin-scale SST Pathfinder vs. 1/12° Pacific HYCOM SST Mean Error




# Comparison of the Zonal Average SST Pathfinder vs. 1/12° Pacific HYCOM



# Velocity Cross-section Across Luzon Strait

Sb-ADCP data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 300 m

Section along 120.75°E between Taiwan and Luzon




Sb-ADCP data from Liang et al. (DSR Pt. II, in press)

6 year mean from HYCOM forced with high-frequency **ECMWF** winds and thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section Across Luzon Strait

Sb-ADCP data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 300 m

Section along 120.75°E between Taiwan and Luzon




Sb-ADCP data from Liang et al. (DSR Pt. II, in press)

6 year mean from HYCOM forced with high-frequency **HR** winds and ECMWF thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section Along Luzon Strait


Sb-ADCP data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 300 m  
Section along 21°N between 118.5°E and 124.0°E



Sb-ADCP data from Liang et al. (DSR Pt. II, in press)  
6 year mean from HYCOM forced with high-frequency **ECMWF** winds and thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section Along Luzon Strait

Sb-ADCP data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 300 m  
Section along 21°N between 118.5°E and 124.0°E




Sb-ADCP data from Liang et al. (DSR Pt. II, in press)  
6 year mean from HYCOM forced with high-frequency **HR** winds and ECMWF thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section East of Taiwan

**Sb-ADCP data (top) vs. 1/12° Pacific HYCOM (bottom) in the upper 300 m**

**Sections at 22°N, 23°N, 24°N and 25°N**




**Sb-ADCP data from Liang et al. (DSR Pt. II, in press)**  
**6 year mean from HYCOM forced with high-frequency ECMWF winds and thermal forcing**  
**No ocean data assimilation in HYCOM**

**Note how the two-core Kuroshio merges to a single jet in both the observations and HYCOM from the south to north along the Taiwan coast**

# Velocity Cross-section East of Taiwan

**Sb-ADCP data (top) vs. 1/12° Pacific HYCOM (bottom) in the upper 300 m**

**Sections at 22°N, 23°N, 24°N and 25°N**





Sb-ADCP data from Liang et al. (DSR Pt. II, in press)

6 year mean from HYCOM forced with high-frequency **HR** winds and ECMWF thermal forcing  
No ocean data assimilation in HYCOM


**Note how the two-core Kuroshio merges to a single jet in both the observations and HYCOM from the south to north along the Taiwan coast**

# Velocity Cross-section at WOCE PCM-1

Current meter data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 1000 m





Note the westward intensification  
of the Kuroshio in the channel  
between Taiwan and the Ryukyu Islands




Current meter data from Lee et al. (2001, JGR)  
6 year mean from HYCOM forced with high-frequency ECMWF winds and thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section at WOCE PCM-1

Current meter data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 1000 m




Note the westward intensification  
of the Kuroshio in the channel  
between Taiwan and the Ryukyu Islands



Current meter data from Lee et al. (2001, JGR)  
6 year mean from HYCOM forced with high-frequency **HR** winds and ECMWF thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section Along the Equator

**TOGA TAO data (top) vs. 1/12° Pacific HYCOM (bottom) in the upper 300 m**  
**Section between 165°E and 110°W**




**TOGA TAO buoy data from Yu and McPhaden (1999, JPO)**  
**6 year mean from HYCOM forced with high-frequency ECMWF winds and thermal forcing**  
**No ocean data assimilation in HYCOM**

# Velocity Cross-section Along the Equator

**TOGA TAO data (top) vs. 1/12° Pacific HYCOM (bottom) in the upper 300 m**


**Section between 165°E and 110°W**



**TOGA TAO buoy data from Yu and McPhaden (1999, JPO)**  
**6 year mean from HYCOM forced with high-frequency **HR** winds and ECMWF thermal forcing**  
**No ocean data assimilation in HYCOM**

# Velocity Cross-section Across the Equator at 135°W

CTD/ADCP data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 400 m  
Section between 8°S and 8°N




CTD/ADCP data from Johnson and McPhaden (2001, JPO)  
6 year mean from HYCOM forced with high-frequency **ECMWF** winds and thermal forcing  
No ocean data assimilation in HYCOM

EUC max velocity:  
90+ cm/s

EUC max velocity:  
80+ cm/s

# Velocity Cross-section Across the Equator at 135°W

CTD/ADCP data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 400 m  
Section between 8°S and 8°N




CTD/ADCP data from Johnson and McPhaden (2001, JPO)  
6 year mean from HYCOM forced with high-frequency **HR** winds and ECMWF thermal forcing  
No ocean data assimilation in HYCOM

EUC max velocity:  
90+ cm/s

EUC max velocity:  
120+ cm/s

# Temperature Cross-section Across the Equator at 135°W


CTD/ADCP data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 400 m  
Section between 8°S and 8°N



CTD/ADCP data from Johnson and McPhaden (2001, JPO)  
6 year mean from HYCOM forced with high-frequency **ECMWF** winds and thermal forcing  
No ocean data assimilation in HYCOM

# Temperature Cross-section Across the Equator at 135°W


CTD/ADCP data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 400 m  
Section between 8°S and 8°N



CTD/ADCP data from Johnson and McPhaden (2001, JPO)  
6 year mean from HYCOM forced with high-frequency **HR** winds and ECMWF thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section Across the Kuroshio at 145°W

Hydrographic data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 1000 m  
Section between 25°N and 50°N




Hydrographic data from Qu et al. (2001, JPO)

6 year mean from HYCOM forced with high-frequency ECMWF winds and thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section Across the Kuroshio at 145°W

Hydrographic data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 1000 m  
Section between 25°N and 50°N




Hydrographic data from Qu et al. (2001, JPO)

6 year mean from HYCOM forced with high-frequency **HR** winds and ECMWF thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section Across the Kuroshio at 155°W

Hydrographic data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 1000 m  
Section between 25°N and 50°N




Hydrographic data from Qu et al. (2001, JPO)

6 year mean from HYCOM forced with high-frequency ECMWF winds and thermal forcing  
No ocean data assimilation in HYCOM

# Velocity Cross-section Across the Kuroshio at 155°W

Hydrographic data (**top**) vs. 1/12° Pacific HYCOM (**bottom**) in the upper 1000 m  
Section between 25°N and 50°N




Hydrographic data from Qu et al. (2001, JPO)

6 year mean from HYCOM forced with high-frequency **HR** winds and ECMWF thermal forcing  
No ocean data assimilation in HYCOM

# Comparison of Currents Around Hawaii

## Composite drifter data vs. 1/12° Pacific HYCOM

Hawaiian Lee  
Counter Current  
(HLCC)




Mean flow field derived  
from 356 WOCE drifters,  
July 1987–March 1998;  
adapted from **Flament  
et al. (1998)** and **Lumpkin  
(1998)**


North Hawaiian  
Ridge Current  
(NHRC)

HYCOM mixed layer current field

HR wind forcing



ECMWF wind forcing




excessively strong HLCC extends all the way to the western boundary




unrealistic NHRC

# Annual Winds Over Hawaii

Wind stress




Wind stress curl



Unrealistic flow around the Hawaiian Islands appears to be related to the anomalously strong wind stress curl dipole in the ECMWF forcing; this is a feature of numerical weather models and not observed ocean wind climatologies

# Linear Response To Wind

## SSH from the linear 1/16°global NRL Layered Ocean Model



Note the unrealistic  
sub-gyre in the southern  
Subtropical Gyre that  
is a linear Sverdrup  
response to the wind  
forcing


# **Methodology to Modify the ECMWF Wind Stress Curl Over the Hawaiian Islands**

1. Define a rectangle in the ECMWF wind stress curl field circumscribing the bull's-eye near Hawaii.
2. Interpolate across the rectangle in both the ECMWF and HR wind stress fields.
3. Subtract the interpolated HR from the pure HR and add the residual to the interpolated ECMWF field.\*
4. Calculate wind stress curl fields and make sure the blending does not create anomalous curl at the rectangle boundaries.
5. Calculate the linear solution using  $1/16^\circ$  global NLOM; if positive results run  $1/12^\circ$  Pacific HYCOM.




\*(Over the Hawaii region the HR stresses are ~40% stronger than ECMWF, so the HR residual is reduced by this amount.)

# Annual Winds Over Hawaii

Wind stress



Wind stress curl




**Successfully reduced magnitude of the wind stress curl dipole over the Hawaiian Islands without introducing anomalous curl at the boundaries**

# Comparison of the Basin-scale Circulation


1/12° Pacific HYCOM: ECMWF winds vs. ECMWF Hawaii modified winds

6-yr mean  
SSH (cm)



ECMWF winds

3-yr mean  
SSH (cm)




ECMWF Hawaii  
modified winds


Note the eastward  
extent of the sub-gyre  
has diminished

# Comparison of the Basin-scale Circulation MODAS climatology vs. 1/12° Pacific HYCOM

Mean dynamic  
height (dyn cm)  
wrt 1000 db



3-yr mean  
SSH (cm)




Forced with high freq. climatological ECMWF winds and a modification around the Hawaiian Islands

# Comparison of Currents Around Hawaii

## Composite drifter data vs. 1/12° Pacific HYCOM

Hawaiian Lee  
Counter Current  
(HLCC)




Mean flow field derived  
from 356 WOCE drifters,  
July 1987–March 1998;  
adapted from Flament  
et al. (1998) and Lumpkin  
(1998)

North Hawaiian  
Ridge Current  
(NHRC)


HYCOM mixed layer current field

ECMWF Hawaii modified wind forcing



more realistic HLCC


ECMWF wind forcing



somewhat improved NHRC

unrealistic NHRC


# Track of Hurricane Juliette




Source: National Hurricane Center

# Evolution of the Coastally Trapped Waves (CTW) Generated By Hurricane Juliette in 1/12° Pacific HYCOM


27 September 2001




28 September 2001




29 September 2001




30 September 2001



1 October 2001



2 October 2001




1/12° Pacific HYCOM forced with FNMOC NOGAPS/HR winds and FNMOC NOGAPS thermal forcing. No data have been assimilated into this model.

↓ Marks the leading edge of the *first* CTW


→ Marks the leading edge of the *second* CTW

# Evolution of the Coastally Trapped Waves (CTW) Generated By Hurricane Juliette in 1/12° Pacific HYCOM


3 October 2001




4 October 2001




5 October 2001




6 October 2001



7 October 2001




8 October 2001



1/12° Pacific HYCOM forced with FNMOC NOGAPS/HR winds and FNMOC NOGAPS thermal forcing. No data have been assimilated into this model.

↓ Marks the leading edge of the *first* CTW

# Observed (solid) vs. Modeled (dashed) Sea Level Along the Mexican Coast Associated With the Coastally Trapped Waves (CTW) Generated by Hurricane Juliette in 2001



1/12° Pacific HYCOM forced with FNMOC NOGAPS/HR winds and FNMOC NOGAPS thermal forcing. No data have been assimilated into this model. Sea level data provided by the University of Hawaii and the Secretaría de Marina de México.